arXiv:quant-ph/0211040v1 8 Nov 2002

Exact Solutions of the Time Dependent Schrödinger Equation in One Space Dimension.

B. Hamprecht
Institut fr Theoretische Physik, Freie Universitt Berlin,
Arnimallee 14, D-14195 Berlin, Germany
e-mails: bodo.hamprecht@physik.fu-berlin.de
(May 16th, 1997)

A closed expression for the harmonic oscillator wave function after the passage of a linear signal with arbitrary time dependence is derived. Transition probabilities are simple to express in terms of Laguerre polynomials. Spontaneous transitions are neglected. The exact result is of some interest for the physics of short laser pulses, since it may serve as an accuracy test for numerical methods.

I. CONSTRUCTION OF THE PROPAGATOR

We consider the Schrödinger Equation:

$$-\frac{\hbar^2}{2m}\frac{\partial^2\Psi(x,t)}{\partial x^2}+\left(\frac{m\omega^2}{2}x^2+xj(t)\right)\Psi(x,t)=i\hbar\frac{\partial\Psi(x,t)}{\partial t} \tag{1.1}$$

The driving force $j(t)$ is supposed to be real and of finite duration. The propagator of the above equation is well known [1].

$$\Psi(x,t,y)=\frac{\alpha}{\sqrt{2\pi i\sin\omega t}}\exp\left\{\frac{i}{\sin\omega t}\left[\alpha^2\left(\frac{x^2+y^2}{2}\cos\omega t-xy\right)+\frac{xx_0+yy_0}{\hbar}+\frac{\chi}{2\alpha^2\hbar^2}\right]\right\} \tag{1.2}$$

where:

$$x_0=-G,\qquad y_0=G\cos\omega t-F\sin\omega t,\qquad chi=G^2\cos\omega t-(FG+2H)\sin\omega t,\qquad \alpha=\sqrt{\frac{m\omega}{\hbar}} \tag{1.3}$$

$$G(t)=\int_0^t dt'j(t')\sin\omega t'\,,\qquad F(t)=\int_0^t dt'j(t')\cos\omega t'$$

$$H(t)=\frac{1}{2}\int_0^t dt'j(t')\int_0^{t'}dt''j(t'')\sin\omega(t''-t')$$

This path integral result describes the propagation of a Gaussian wave packet, which starts from δ-function shape at time $t=0$. More easily it may be derived by solving the time dependent Schrödinger equation (1.1).

We notice that the wave packet

$$\Psi(x,t)=\exp\left\{-A(t)\frac{\alpha^2x^2}{2}+iB(t)\alpha x-\frac{C(t)}{2}\right\} \tag{1.4}$$

solves equation (1.1), if the complex valued functions $A(t),B(t)$ and $C(t)$ are solutions of the following set of ordinary differential equations:

$$\frac{d}{dt}A=i\omega(1-A^2)\qquad \frac{d}{dt}B=-i\omega A\,B-\frac{j(t)}{\hbar\alpha}\qquad \frac{d}{dt}C=i\omega(A+B^2) \tag{1.5}$$

Since we want to have a δ-function initially:

$$\Psi(x,0,y):=\Psi(x,0)=\delta(x-y)=\lim_{\tau\to 0}\frac{1}{\sqrt{\pi\tau}}\exp\left(-\frac{(x-y)^2}{\tau}\right)$$

A,B and C have to be solutions of (1.5) with a singularity at $t=0$:

$$A(t) = -i \cot \omega t$$
$$B(t) = -\frac{\frac{G(t)}{\hbar} + \alpha^2 y}{\alpha \sin \omega t}$$
$$C(t) = \log(2\pi i \sin \omega t) - i\alpha^2 y^2 \cot \omega t - i \left(2y + \frac{G}{\alpha^2 \hbar}\right) \frac{G \cot \omega t - F}{\hbar} + \frac{2i}{\alpha^2 \hbar^2} H \tag{1.6}$$

Substituting eqs. (1.6) in equ. (1.4) yields the Propagator (1.2).

II. TRANSITION PROBABILITIES

Transition probability amplitudes between energy eigenstates of the harmonic oscillator are given by:

$$\begin{aligned} a_{m,n} &= \int\limits_{-\infty}^{\infty} \psi_m^*(x,t)\Psi(x,t,y)\psi_n(y,0)dx\,dy \\ &= \alpha \int\limits_{-\infty}^{\infty} \frac{H_m(\alpha x)\Psi(x,y,t)H_n(\alpha y)}{\sqrt{2^{n+m} m! n! \pi}} \exp\left\{-\frac{x^2+y^2}{2}\alpha^2 + i\hbar\omega t\left(m + \frac{1}{2}\right)\right\} dx\,dy \end{aligned} \tag{2.1}$$

This integral is evaluated making use of the generating function $\exp(2zx - z^2)$ for the Hermite polynomials $H_n(x)$ (cf. Appendix), yielding for $m \leq n$:

$$a_{n,m} = \sqrt{\frac{m!}{n!}} \exp\left\{-\frac{|r|^2}{2} - \frac{iH}{\alpha^2 \hbar^2}\right\} (-ir^*)^{n-m} L_m^{(n-m)}(|r|^2) \tag{2.2}$$

where $L_m^{(n-m)}(x)$ is a generalised Laguerre polynomial

and

$$r = \frac{F - iG}{\sqrt{2}\alpha\hbar} = \frac{1}{\sqrt{2}\alpha\hbar} \int\limits_0^t dt' j(t') e^{i\omega t'}$$

Equation ([**?**]) is our main result.
For transitions from the ground state, i.e. $m = 0$, equation (8) reduces to the simple expression:

$$\begin{aligned} a_n &= \frac{1}{\sqrt{n!}} \exp\left\{-\frac{|r|^2 + iH}{2}\right\} (-ir^*)^n \\ |a_n|^2 &= \frac{R^n}{n!} e^{-R} \qquad \text{where} \qquad R = |r|^2 \end{aligned} \tag{2.3}$$

III. EXITATION OF WAVE PACKETS

If the pulse $j(t)$ is applied to the ground state of the harmonic oscillator, it will produce a wave packet, which is easily determined, applying the propagator to the wave function of the ground state:

$$\Psi(x,t) = \sqrt[4]{\frac{\alpha^2}{\pi}} \int\limits_{-\infty}^{\infty} \Psi(x,t,y) \exp\left(-\frac{\alpha^2 y^2}{2}\right) dy \tag{3.1}$$

The integral evaluates to:

$$\Psi(x,t) = \sqrt[4]{\frac{\alpha^2}{\pi}} \exp\left[-\frac{\alpha^2}{2}\left(x - x_0)^2\right) - \chi(t)\right] \tag{3.2}$$

where

$$x_0 = -i\frac{(F+iG)e^{-i\omega t}}{\hbar\alpha^2}$$
$$\chi(t) = \frac{(F+iG)(F\cos\omega t + G\sin\omega t)e^{-i\omega t} + 2iH}{2\alpha^2\hbar^2} + \frac{1}{2}\, i\omega t$$

describing an oscillating Gaussian wave packet of constant width $\Delta x^2 = 1/(2\alpha^2)$.
The expectation values for position and momentum turn out to be:

$$\langle x\rangle = \frac{G\cos\omega t - F\sin\omega t}{\alpha^2\hbar} \qquad \langle p\rangle = -(F\cos\omega t + G\sin\omega t) \tag{3.3}$$

IV. APPENDIX

We evaluate equation (2.1), using the generating function for the Hermite polynomials. If $\gamma_{m,n}$ is the coefficient of $w^m z^n$ in the Taylor expansion of:

$$J = \alpha \int_{-\infty}^{\infty} \Psi(x,t,y) e^{2(wx+zy)\alpha - w^2 - z^2 - \frac{x^2+y^2}{2}\alpha^2}\, dx\, dy \qquad \text{then}$$

$$a_{m,n} = \sqrt{\frac{n!m!}{2^{m+n}\pi}} \exp\left\{ i\hbar\omega t\left(m + \frac{1}{2}\right)\right\} \gamma_{m,n} \tag{4.1}$$

Now:

$$\begin{aligned} J &= \frac{1}{\sqrt{2\pi i \sin t}} \int_{-\infty}^{\infty} \exp\left\{-X^T A X + 2P^T X - Q\right\} \alpha^2 dX \\ &= \sqrt{\frac{\pi}{2i\sin t\, det A}} \exp\left\{P^T A^{-1} P - Q\right\} \end{aligned} \tag{4.2}$$

where:

$$X = \alpha\begin{pmatrix} x \\ y\end{pmatrix} \qquad P = \begin{pmatrix} w - i\frac{G}{2\alpha\hbar\sin\omega t} \\ z - \frac{i}{2\alpha\hbar}(F - G\cot\omega t)\end{pmatrix} \qquad A = \begin{pmatrix} \frac{1}{2} - \frac{i}{2}\cot\omega t & \frac{i}{2\sin\omega t} \\ \frac{i}{2\sin\omega t} & \frac{1}{2} - \frac{i}{2}\cot\omega t \end{pmatrix}$$
$$Q = w^2 + z^2 + \frac{i}{2\alpha^2\hbar^2}(F - G\cot\omega t)G + \frac{i}{\alpha^2\hbar^2}H$$

We find:

$$det\, A = -i\frac{e^{i\omega t}}{2\sin\omega t} \qquad A^{-1} = \begin{pmatrix} 1 & e^{-i\omega t} \\ e^{-i\omega t} & 1 \end{pmatrix} \tag{4.3}$$

$$P^T A^{-1} P - Q = 2wz\, e^{-i\omega t} - iz\frac{F - iG}{\alpha\hbar} - iw\frac{F+iG}{\alpha\hbar}e^{-i\omega t} - \frac{F^2 + G^2 + 4iH}{4\alpha^2\hbar^2} \tag{4.4}$$

Therefore:

$$a_{m,n} = \sqrt{\frac{n!m!}{2^{n+m}}} \exp\left\{-\frac{F^2+G^2+4iH}{4\alpha^2\hbar^2}\right\} \sum_{k=0}^{m} \frac{\left(2e^{-i\omega t}\right)^k \left(\frac{F+iG}{\alpha\hbar}\right)^{m-k} \left(\frac{F-iG}{\alpha\hbar}\right)^{n-k} (-i)^{m+n-2k} e^{ik\omega t}}{k!(m-k)!(n-k)!}$$

Extracting common factors from the sum and replacing k by $m-k$ in the summation, we obtain:

$$a_{n.m} = \sqrt{n!m!} \exp\left\{-\frac{F^2+G^2+4iH}{4\alpha^2\hbar^2}\right\} \left(-i\frac{F-iG}{\sqrt{2}\alpha\hbar}\right)^{n-m} \sum_{k=0}^{m} \frac{\left(-\frac{F^2+G^2}{2\alpha^2\hbar^2}\right)^k}{k!(m-k)!(n-m+k)!}$$

The finite sum in this expression defines a Laguerre polynomial. Therefore:

$$a_{n.m} = \sqrt{\frac{m!}{n!}} \exp\left\{-\frac{|r|^2}{2} - \frac{iH}{\alpha^2\hbar^2}\right\} (-ir^*)^{n-m} L_m^{(n-m)}(|r|^2) \tag{4.5}$$

where:

$$r = \frac{F + iG}{\sqrt{2}\alpha\hbar} = \frac{1}{\sqrt{2}\alpha\hbar}\int\limits_0^t dt' j(t') e^{i\omega t'}$$

which proves equation (2.2).

---

[1] Hagen Kleinert: **Path Integrals in Quantum Mechanics, Statistics and Polymer Physics**, 2^{nd} edition, World Scientific(1995).